

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Average	97.5%
Colorectal Cancer Screening		Above Expected	74.9%
Optimal Asthma Control - Adults		Expected	58.2%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	47.5%
HbA1c Control		Expected	74.7%
BP Control		Expected	81.4%
Daily Aspirin Use		Expected	98.5%
Statin Use		Expected	86.3%
Tobacco-Free		Above Expected	90.6%
Optimal Vascular Care		Above Expected	62.0%
BP Control		Expected	83.4%
Daily Aspirin Use		Expected	86.3%
Statin Use		Expected	92.3%
Tobacco-Free		Above Expected	92.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).