

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	59.8%
	Follow-up at Six Months	Expected	40.9%
	Response at Six Months	Expected	12.3%
	Remission at Six Months	Below Expected	4.4%
	Follow-up at 12 Months	Below Expected	34.5%
	Response at 12 Months	Below Expected	6.4%
	Remission at 12 Months	Below Expected	4.4%
Adolescent Mental Health and/or Depression Screening*		Below	66.5%
Colorectal Cancer Screening		Below Expected	60.0%
Optimal Asthma Control - Adults		Below Expected	21.9%
Optimal Asthma Control - Children		Below Expected	16.9%
Optimal Diabetes Care		Below Expected	36.5%
HbA1c Control		Expected	68.4%
BP Control		Above Expected	89.4%
Daily Aspirin Use		Below Expected	92.7%
Statin Use		Below Expected	78.5%
Tobacco-Free		Below Expected	79.4%
Optimal Vascular Care		Below Expected	31.9%
BP Control		Expected	84.4%
Daily Aspirin Use		Below Expected	53.5%
Statin Use		Below Expected	83.9%
Tobacco-Free		Below Expected	75.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).