

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	55.9%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Below	84.9%
Colorectal Cancer Screening		Expected	66.9%
Optimal Asthma Control - Adults		Expected	44.8%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	38.7%
HbA1c Control		Expected	71.4%
BP Control		Expected	85.7%
Daily Aspirin Use		Below Expected	94.5%
Statin Use		Expected	80.3%
Tobacco-Free		Below Expected	76.5%
Optimal Vascular Care		Expected	40.2%
BP Control		Expected	90.2%
Daily Aspirin Use		Below Expected	62.8%
Statin Use		Expected	84.3%
Tobacco-Free		Expected	73.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).