

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	61.8%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Average	92.3%
Colorectal Cancer Screening		Below Expected	61.1%
Optimal Asthma Control - Adults		Below Expected	18.5%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	36.8%
HbA1c Control		Expected	70.9%
BP Control		Above Expected	89.6%
Daily Aspirin Use		Expected	96.7%
Statin Use		Expected	81.9%
Tobacco-Free		Below Expected	75.8%
Optimal Vascular Care		Below Expected	40.0%
BP Control		Above Expected	94.0%
Daily Aspirin Use		Below Expected	74.0%
Statin Use		Below Expected	79.0%
Tobacco-Free		Expected	71.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).