

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Above	97.8%
Colorectal Cancer Screening		Expected	72.2%
Optimal Asthma Control - Adults		Below Expected	40.0%
Optimal Asthma Control - Children		Expected	59.5%
Optimal Diabetes Care		Expected	49.5%
HbA1c Control		Expected	76.3%
BP Control		Above Expected	86.2%
Daily Aspirin Use		Expected	98.9%
Statin Use		Below Expected	85.2%
Tobacco-Free		Expected	87.6%
Optimal Vascular Care		Above Expected	63.7%
BP Control		Expected	84.4%
Daily Aspirin Use		Above Expected	89.8%
Statin Use		Expected	93.5%
Tobacco-Free		Expected	87.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).