

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Above	97.2%
Colorectal Cancer Screening		Expected	70.9%
Optimal Asthma Control - Adults		Below Expected	45.4%
Optimal Asthma Control - Children		Below Expected	32.2%
Optimal Diabetes Care		Above Expected	52.1%
HbA1c Control		Above Expected	78.2%
BP Control		Above Expected	85.1%
Daily Aspirin Use		Expected	99.0%
Statin Use		Above Expected	91.4%
Tobacco-Free		Expected	83.0%
Optimal Vascular Care		Expected	57.1%
BP Control		Expected	82.4%
Daily Aspirin Use		Expected	89.8%
Statin Use		Above Expected	95.0%
Tobacco-Free		Expected	77.6%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).