

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Above	98.8%
Colorectal Cancer Screening		Below Expected	68.4%
Optimal Asthma Control - Adults		Expected	48.0%
Optimal Asthma Control - Children		Below Expected	30.0%
Optimal Diabetes Care		Expected	46.6%
HbA1c Control		Expected	75.4%
BP Control		Expected	86.2%
Daily Aspirin Use		Expected	99.7%
Statin Use		Expected	85.9%
Tobacco-Free		Expected	84.7%
Optimal Vascular Care		Expected	57.1%
BP Control		Expected	82.5%
Daily Aspirin Use		Expected	91.6%
Statin Use		Expected	96.1%
Tobacco-Free		Expected	77.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).