

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	86.9%
	Follow-up at Six Months	Expected	55.9%
	Response at Six Months	Expected	19.9%
	Remission at Six Months	Expected	12.9%
	Follow-up at 12 Months	Expected	59.0%
	Response at 12 Months	Expected	24.6%
	Remission at 12 Months	Expected	13.3%
Adolescent Mental Health and/or Depression Screening*		Above	99.4%
Colorectal Cancer Screening		Below Expected	69.4%
Optimal Asthma Control - Adults		Expected	50.6%
Optimal Asthma Control - Children		Below Expected	28.7%
Optimal Diabetes Care		Below Expected	39.0%
HbA1c Control		Expected	71.1%
BP Control		Below Expected	75.7%
Daily Aspirin Use		Expected	98.7%
Statin Use		Below Expected	85.9%
Tobacco-Free		Below Expected	81.5%
Optimal Vascular Care		Below Expected	46.0%
BP Control		Below Expected	74.3%
Daily Aspirin Use		Expected	88.0%
Statin Use		Below Expected	88.3%
Tobacco-Free		Expected	78.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).