

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Below	88.9%
Colorectal Cancer Screening		Above Expected	78.4%
Optimal Asthma Control - Adults		Below Expected	30.1%
Optimal Asthma Control - Children		Below Expected	26.1%
Optimal Diabetes Care		Expected	51.5%
HbA1c Control		Expected	74.1%
BP Control		Expected	81.4%
Daily Aspirin Use		Above Expected	99.5%
Statin Use		Expected	89.8%
Tobacco-Free		Above Expected	90.3%
Optimal Vascular Care		Expected	61.8%
BP Control		Expected	82.8%
Daily Aspirin Use		Expected	88.6%
Statin Use		Expected	93.2%
Tobacco-Free		Above Expected	88.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).