

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	80.7%
	Follow-up at Six Months	Expected	53.0%
	Response at Six Months	Expected	25.5%
	Remission at Six Months	Expected	12.1%
	Follow-up at 12 Months	Expected	47.0%
	Response at 12 Months	Expected	20.1%
	Remission at 12 Months	Expected	13.4%
Adolescent Mental Health and/or Depression Screening*		Below	82.7%
Colorectal Cancer Screening		Expected	67.2%
Optimal Asthma Control - Adults		Above Expected	69.0%
Optimal Asthma Control - Children		Expected	64.4%
Optimal Diabetes Care		Expected	48.5%
HbA1c Control		Expected	73.9%
BP Control		Expected	80.3%
Daily Aspirin Use		Expected	99.5%
Statin Use		Expected	87.6%
Tobacco-Free		Expected	87.0%
Optimal Vascular Care		Expected	58.3%
BP Control		Expected	82.1%
Daily Aspirin Use		Expected	85.7%
Statin Use		Expected	89.3%
Tobacco-Free		Expected	86.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).