

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	85.9%
	Follow-up at Six Months	Expected	33.8%
	Response at Six Months	Expected	8.8%
	Remission at Six Months	Expected	6.3%
	Follow-up at 12 Months	Expected	40.0%
	Response at 12 Months	Expected	10.0%
	Remission at 12 Months	Expected	5.0%
Adult Depression	PHQ-9 Utilization*	Below	73.0%
	Follow-up at Six Months	Expected	50.0%
	Response at Six Months	Expected	20.7%
	Remission at Six Months	Expected	13.9%
	Follow-up at 12 Months	Expected	53.5%
	Response at 12 Months	Expected	22.3%
	Remission at 12 Months	Expected	13.7%
Adolescent Mental Health and/or Depression Screening*		Below	81.8%
Colorectal Cancer Screening		Above Expected	75.8%
Optimal Asthma Control - Adults		Expected	56.5%
Optimal Asthma Control - Children		Expected	47.0%
Optimal Diabetes Care		Expected	50.9%
HbA1c Control		Expected	76.5%
BP Control		Expected	79.4%
Daily Aspirin Use		Expected	99.1%
Statin Use		Above Expected	94.2%
Tobacco-Free		Expected	84.8%
Optimal Vascular Care		Expected	53.4%
BP Control		Below Expected	74.8%
Daily Aspirin Use		Above Expected	90.8%
Statin Use		Expected	92.2%
Tobacco-Free		Expected	81.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).