

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	31.9%
	Response at Six Months	Expected	8.5%
	Remission at Six Months	Expected	6.4%
	Follow-up at 12 Months	Expected	38.3%
	Response at 12 Months	Expected	8.5%
	Remission at 12 Months	Expected	6.4%
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Below Expected	21.2%
	Response at Six Months	Expected	12.1%
	Remission at Six Months	Expected	6.1%
	Follow-up at 12 Months	Expected	27.3%
	Response at 12 Months	Expected	18.2%
	Remission at 12 Months	Expected	9.1%
Adolescent Mental Health and/or Depression Screening*		Above	99.1%
Colorectal Cancer Screening		-	-
Optimal Asthma Control - Adults		Expected	48.2%
Optimal Asthma Control - Children		Expected	53.1%
Optimal Diabetes Care		-	-
HbA1c Control		-	-
BP Control		-	-
Daily Aspirin Use		-	-
Statin Use		-	-
Tobacco-Free		-	-
Optimal Vascular Care		-	-
BP Control		-	-
Daily Aspirin Use		-	-
Statin Use		-	-
Tobacco-Free		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).