

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Above	97.1%
Colorectal Cancer Screening		Above Expected	77.7%
Optimal Asthma Control - Adults		Expected	50.9%
Optimal Asthma Control - Children		Below Expected	42.2%
Optimal Diabetes Care		Above Expected	57.6%
HbA1c Control		Above Expected	82.9%
BP Control		Expected	83.3%
Daily Aspirin Use		Expected	98.6%
Statin Use		Expected	89.7%
Tobacco-Free		Above Expected	91.8%
Optimal Vascular Care		Above Expected	63.2%
BP Control		Expected	82.3%
Daily Aspirin Use		Expected	86.0%
Statin Use		Expected	93.6%
Tobacco-Free		Above Expected	92.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).