

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	94.3%
	Follow-up at Six Months	Expected	59.7%
	Response at Six Months	Expected	19.5%
	Remission at Six Months	Expected	9.1%
	Follow-up at 12 Months	Expected	55.8%
	Response at 12 Months	Expected	24.7%
	Remission at 12 Months	Expected	16.9%
Adolescent Mental Health and/or Depression Screening*		Below	85.2%
Colorectal Cancer Screening		Below Expected	68.0%
Optimal Asthma Control - Adults		Above Expected	77.3%
Optimal Asthma Control - Children		Expected	72.1%
Optimal Diabetes Care		Expected	50.2%
HbA1c Control		Expected	75.3%
BP Control		Expected	82.4%
Daily Aspirin Use		Expected	99.7%
Statin Use		Expected	90.9%
Tobacco-Free		Expected	88.1%
Optimal Vascular Care		Expected	59.3%
BP Control		Expected	85.2%
Daily Aspirin Use		Expected	86.1%
Statin Use		Expected	95.4%
Tobacco-Free		Expected	84.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).