

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	84.8%
	Follow-up at Six Months	Expected	56.3%
	Response at Six Months	Expected	18.8%
	Remission at Six Months	Expected	9.4%
	Follow-up at 12 Months	Expected	53.1%
	Response at 12 Months	Expected	21.9%
	Remission at 12 Months	Expected	9.4%
Adult Depression	PHQ-9 Utilization*	Above	88.9%
	Follow-up at Six Months	Above Expected	68.2%
	Response at Six Months	Expected	26.9%
	Remission at Six Months	Expected	13.9%
	Follow-up at 12 Months	Above Expected	66.8%
	Response at 12 Months	Expected	24.2%
	Remission at 12 Months	Expected	11.7%
Adolescent Mental Health and/or Depression Screening*		Below	86.3%
Colorectal Cancer Screening		Expected	71.2%
Optimal Asthma Control - Adults		Above Expected	63.0%
Optimal Asthma Control - Children		Expected	66.7%
Optimal Diabetes Care		Expected	49.6%
HbA1c Control		Above Expected	80.4%
BP Control		Expected	78.9%
Daily Aspirin Use		Expected	98.8%
Statin Use		Expected	89.0%
Tobacco-Free		Below Expected	83.9%
Optimal Vascular Care		Expected	58.5%
BP Control		Expected	81.9%
Daily Aspirin Use		Above Expected	90.0%
Statin Use		Expected	93.9%
Tobacco-Free		Expected	81.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).