

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Above	98.9%
Colorectal Cancer Screening		Expected	75.1%
Optimal Asthma Control - Adults		Below Expected	40.4%
Optimal Asthma Control - Children		Expected	40.0%
Optimal Diabetes Care		Expected	51.5%
HbA1c Control		Expected	77.4%
BP Control		Above Expected	88.0%
Daily Aspirin Use		Expected	99.2%
Statin Use		Below Expected	83.2%
Tobacco-Free		Expected	90.4%
Optimal Vascular Care		Above Expected	65.3%
BP Control		Expected	85.9%
Daily Aspirin Use		Expected	89.6%
Statin Use		Expected	92.8%
Tobacco-Free		Expected	88.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).