

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	39.1%
	Response at Six Months	Expected	14.5%
	Remission at Six Months	Expected	5.8%
	Follow-up at 12 Months	Expected	36.2%
	Response at 12 Months	Expected	17.4%
	Remission at 12 Months	Expected	4.4%
Adolescent Mental Health and/or Depression Screening*		Above	100.0%
Colorectal Cancer Screening		Above Expected	77.6%
Optimal Asthma Control - Adults		Above Expected	70.3%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	48.8%
HbA1c Control		Expected	79.8%
BP Control		Expected	79.0%
Daily Aspirin Use		Expected	99.2%
Statin Use		Expected	92.7%
Tobacco-Free		Expected	81.1%
Optimal Vascular Care		Expected	53.2%
BP Control		Expected	84.8%
Daily Aspirin Use		Expected	88.6%
Statin Use		Expected	93.0%
Tobacco-Free		Below Expected	71.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).