

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Below Expected	31.7%
	Response at Six Months	Expected	14.1%
	Remission at Six Months	Expected	9.2%
	Follow-up at 12 Months	Below Expected	31.0%
	Response at 12 Months	Below Expected	10.6%
	Remission at 12 Months	Below Expected	4.2%
Adolescent Mental Health and/or Depression Screening*		Above	98.1%
Colorectal Cancer Screening		Above Expected	76.8%
Optimal Asthma Control - Adults		Expected	60.2%
Optimal Asthma Control - Children		Expected	57.1%
Optimal Diabetes Care		Expected	55.2%
HbA1c Control		Expected	78.5%
BP Control		Expected	87.2%
Daily Aspirin Use		Expected	99.1%
Statin Use		Expected	92.5%
Tobacco-Free		Expected	81.5%
Optimal Vascular Care		Expected	58.8%
BP Control		Above Expected	89.5%
Daily Aspirin Use		Expected	86.4%
Statin Use		Expected	91.5%
Tobacco-Free		Expected	77.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).