

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	75.0%
	Follow-up at Six Months	Expected	43.5%
	Response at Six Months	Expected	13.0%
	Remission at Six Months	Expected	4.4%
	Follow-up at 12 Months	Expected	28.3%
	Response at 12 Months	Expected	4.4%
	Remission at 12 Months	Expected	4.4%
Adult Depression	PHQ-9 Utilization*	Above	88.1%
	Follow-up at Six Months	Expected	48.8%
	Response at Six Months	Expected	17.2%
	Remission at Six Months	Expected	11.4%
	Follow-up at 12 Months	Below Expected	44.2%
	Response at 12 Months	Expected	17.2%
	Remission at 12 Months	Expected	10.4%
Adolescent Mental Health and/or Depression Screening*		Below	89.8%
Colorectal Cancer Screening		Expected	72.4%
Optimal Asthma Control - Adults		Above Expected	63.4%
Optimal Asthma Control - Children		Expected	55.6%
Optimal Diabetes Care		Above Expected	53.5%
HbA1c Control		Above Expected	80.2%
BP Control		Above Expected	85.4%
Daily Aspirin Use		Expected	98.8%
Statin Use		Expected	89.0%
Tobacco-Free		Expected	85.4%
Optimal Vascular Care		Expected	58.6%
BP Control		Expected	86.2%
Daily Aspirin Use		Expected	88.9%
Statin Use		Expected	93.4%
Tobacco-Free		Expected	81.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).