

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	Expected	47.5%
	Response at Six Months	Expected	15.0%
	Remission at Six Months	Expected	7.5%
	Follow-up at 12 Months	Expected	42.5%
	Response at 12 Months	Expected	15.0%
	Remission at 12 Months	Expected	7.5%
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	45.2%
	Response at Six Months	Below Expected	10.1%
	Remission at Six Months	Expected	5.4%
	Follow-up at 12 Months	Expected	41.1%
	Response at 12 Months	Expected	14.3%
	Remission at 12 Months	Expected	7.1%
Adolescent Mental Health and/or Depression Screening*		Above	98.8%
Colorectal Cancer Screening		Above Expected	79.9%
Optimal Asthma Control - Adults		Above Expected	65.2%
Optimal Asthma Control - Children		Above Expected	71.4%
Optimal Diabetes Care		Expected	52.2%
HbA1c Control		Above Expected	81.6%
BP Control		Expected	82.4%
Daily Aspirin Use		Expected	98.5%
Statin Use		Expected	86.3%
Tobacco-Free		Expected	83.3%
Optimal Vascular Care		Expected	57.6%
BP Control		Expected	79.7%
Daily Aspirin Use		Expected	83.8%
Statin Use		Expected	91.1%
Tobacco-Free		Expected	83.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).