

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	42.1%
	Response at Six Months	Expected	13.8%
	Remission at Six Months	Expected	7.6%
	Follow-up at 12 Months	Expected	39.0%
	Response at 12 Months	Expected	13.2%
	Remission at 12 Months	Expected	5.7%
Adolescent Mental Health and/or Depression Screening*		Above	97.4%
Colorectal Cancer Screening		Expected	70.2%
Optimal Asthma Control - Adults		Expected	55.7%
Optimal Asthma Control - Children		Expected	50.8%
Optimal Diabetes Care		Expected	45.7%
HbA1c Control		Expected	76.6%
BP Control		Expected	77.1%
Daily Aspirin Use		Expected	98.7%
Statin Use		Expected	91.3%
Tobacco-Free		Below Expected	78.2%
Optimal Vascular Care		Expected	48.0%
BP Control		Expected	77.0%
Daily Aspirin Use		Expected	89.5%
Statin Use		Expected	94.5%
Tobacco-Free		Below Expected	70.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).