

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	71.6%
	Follow-up at Six Months	Expected	39.3%
	Response at Six Months	Below Expected	6.5%
	Remission at Six Months	Expected	3.7%
	Follow-up at 12 Months	Below Expected	32.7%
	Response at 12 Months	Below Expected	5.6%
	Remission at 12 Months	Below Expected	1.9%
Adolescent Mental Health and/or Depression Screening*		Below	81.8%
Colorectal Cancer Screening		Below Expected	45.1%
Optimal Asthma Control - Adults		Below Expected	25.3%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Below Expected	30.7%
HbA1c Control		Expected	67.0%
BP Control		Below Expected	69.3%
Daily Aspirin Use		Above Expected	100.0%
Statin Use		Expected	85.4%
Tobacco-Free		Below Expected	71.2%
Optimal Vascular Care		Expected	43.4%
BP Control		Expected	68.4%
Daily Aspirin Use		Expected	92.1%
Statin Use		Expected	97.4%
Tobacco-Free		Below Expected	63.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).