

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	81.7%
	Follow-up at Six Months	Below Expected	19.1%
	Response at Six Months	Below Expected	8.6%
	Remission at Six Months	Below Expected	3.8%
	Follow-up at 12 Months	Below Expected	25.7%
	Response at 12 Months	Expected	13.3%
	Remission at 12 Months	Expected	9.5%
Adolescent Mental Health and/or Depression Screening*		Average	95.2%
Colorectal Cancer Screening		Below Expected	67.9%
Optimal Asthma Control - Adults		Below Expected	2.7%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	48.4%
HbA1c Control		Expected	76.5%
BP Control		Expected	82.3%
Daily Aspirin Use		Expected	97.6%
Statin Use		Expected	87.6%
Tobacco-Free		Expected	87.0%
Optimal Vascular Care		Expected	58.3%
BP Control		Expected	86.1%
Daily Aspirin Use		Below Expected	74.7%
Statin Use		Expected	93.3%
Tobacco-Free		Above Expected	89.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).