

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	91.7%
	Follow-up at Six Months	Expected	39.2%
	Response at Six Months	Expected	10.3%
	Remission at Six Months	Expected	4.1%
	Follow-up at 12 Months	Below Expected	28.9%
	Response at 12 Months	Expected	12.4%
	Remission at 12 Months	Expected	4.1%
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Expected	58.9%
Optimal Asthma Control - Adults		Below Expected	17.7%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	39.6%
HbA1c Control		Expected	68.4%
BP Control		Above Expected	89.9%
Daily Aspirin Use		Expected	99.3%
Statin Use		Expected	84.9%
Tobacco-Free		Expected	75.5%
Optimal Vascular Care		Expected	52.1%
BP Control		Above Expected	95.8%
Daily Aspirin Use		Expected	85.4%
Statin Use		Expected	85.4%
Tobacco-Free		Expected	72.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).