

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	76.6%
	Follow-up at Six Months	Below Expected	32.6%
	Response at Six Months	Below Expected	12.1%
	Remission at Six Months	Below Expected	6.2%
	Follow-up at 12 Months	Below Expected	30.5%
	Response at 12 Months	Below Expected	8.8%
	Remission at 12 Months	Below Expected	4.3%
Adolescent Mental Health and/or Depression Screening*		Below	84.9%
Colorectal Cancer Screening		Above Expected	76.2%
Optimal Asthma Control - Adults		Below Expected	38.0%
Optimal Asthma Control - Children		Below Expected	30.0%
Optimal Diabetes Care		Below Expected	41.3%
HbA1c Control		Expected	70.0%
BP Control		Expected	84.9%
Daily Aspirin Use		Below Expected	94.9%
Statin Use		Expected	83.8%
Tobacco-Free		Expected	83.0%
Optimal Vascular Care		Below Expected	37.8%
BP Control		Above Expected	88.6%
Daily Aspirin Use		Below Expected	60.0%
Statin Use		Below Expected	83.3%
Tobacco-Free		Expected	79.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).