

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	42.4%
	Response at Six Months	Expected	13.5%
	Remission at Six Months	Expected	6.7%
	Follow-up at 12 Months	Below Expected	35.7%
	Response at 12 Months	Below Expected	10.4%
	Remission at 12 Months	Expected	5.1%
Adolescent Mental Health and/or Depression Screening*		Above	98.4%
Colorectal Cancer Screening		Expected	59.3%
Optimal Asthma Control - Adults		Expected	50.7%
Optimal Asthma Control - Children		Expected	41.5%
Optimal Diabetes Care		Expected	40.9%
HbA1c Control		Expected	69.8%
BP Control		Expected	79.9%
Daily Aspirin Use		Expected	99.0%
Statin Use		Above Expected	91.2%
Tobacco-Free		Expected	79.7%
Optimal Vascular Care		Expected	45.8%
BP Control		Expected	75.0%
Daily Aspirin Use		Expected	88.9%
Statin Use		Above Expected	94.8%
Tobacco-Free		Below Expected	66.7%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).