

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	59.3%
	Follow-up at Six Months	Above Expected	80.3%
	Response at Six Months	Above Expected	42.0%
	Remission at Six Months	Above Expected	22.9%
	Follow-up at 12 Months	Above Expected	72.6%
	Response at 12 Months	Above Expected	42.7%
	Remission at 12 Months	Above Expected	27.4%
Adolescent Mental Health and/or Depression Screening*		Average	93.8%
Colorectal Cancer Screening		Expected	72.9%
Optimal Asthma Control - Adults		Expected	46.6%
Optimal Asthma Control - Children		Expected	68.9%
Optimal Diabetes Care		Expected	46.2%
HbA1c Control		Expected	73.1%
BP Control		Expected	83.4%
Daily Aspirin Use		Expected	99.0%
Statin Use		Above Expected	91.7%
Tobacco-Free		Below Expected	80.6%
Optimal Vascular Care		Expected	58.1%
BP Control		Expected	82.3%
Daily Aspirin Use		Above Expected	90.6%
Statin Use		Expected	94.6%
Tobacco-Free		Expected	80.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).