

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		Expected	73.7%
Optimal Asthma Control - Adults		-	-
Optimal Asthma Control - Children		-	-
<b>Optimal Diabetes Care</b>		<b>Expected</b>	<b>45.7%</b>
<i>HbA1c Control</i>		<i>Expected</i>	78.3%
<i>BP Control</i>		<i>Expected</i>	87.0%
<i>Daily Aspirin Use</i>		<i>Expected</i>	98.9%
<i>Statin Use</i>		<i>Expected</i>	90.2%
<i>Tobacco-Free</i>		<i>Expected</i>	73.9%
<b>Optimal Vascular Care</b>		<b>Expected</b>	<b>57.1%</b>
<i>BP Control</i>		<i>Expected</i>	82.9%
<i>Daily Aspirin Use</i>		<i>Expected</i>	85.7%
<i>Statin Use</i>		<i>Expected</i>	91.4%
<i>Tobacco-Free</i>		<i>Expected</i>	74.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).