

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	66.2%
	Follow-up at Six Months	Above Expected	90.9%
	Response at Six Months	Above Expected	54.6%
	Remission at Six Months	Above Expected	33.3%
	Follow-up at 12 Months	Above Expected	81.8%
	Response at 12 Months	Above Expected	51.5%
	Remission at 12 Months	Above Expected	48.5%
Adolescent Mental Health and/or Depression Screening*		Average	100.0%
Colorectal Cancer Screening		Above Expected	79.8%
Optimal Asthma Control - Adults		Expected	61.9%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Above Expected	62.8%
HbA1c Control		Expected	82.9%
BP Control		Expected	87.8%
Daily Aspirin Use		Expected	99.4%
Statin Use		Expected	95.1%
Tobacco-Free		Expected	86.0%
Optimal Vascular Care		Expected	68.0%
BP Control		Expected	87.8%
Daily Aspirin Use		Expected	91.8%
Statin Use		Expected	96.6%
Tobacco-Free		Expected	87.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).