

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	81.5%
	Follow-up at Six Months	Expected	48.6%
	Response at Six Months	Expected	11.4%
	Remission at Six Months	Expected	5.7%
	Follow-up at 12 Months	Expected	48.6%
	Response at 12 Months	Expected	25.7%
	Remission at 12 Months	Expected	11.4%
Adult Depression	PHQ-9 Utilization*	Below	57.3%
	Follow-up at Six Months	Above Expected	75.4%
	Response at Six Months	Above Expected	32.7%
	Remission at Six Months	Above Expected	22.2%
	Follow-up at 12 Months	Above Expected	68.2%
	Response at 12 Months	Above Expected	35.4%
	Remission at 12 Months	Above Expected	21.3%
Adolescent Mental Health and/or Depression Screening*		Above	96.4%
Colorectal Cancer Screening		Above Expected	76.2%
Optimal Asthma Control - Adults		Expected	53.9%
Optimal Asthma Control - Children		Expected	50.7%
Optimal Diabetes Care		Expected	48.5%
HbA1c Control		Expected	74.8%
BP Control		Expected	83.6%
Daily Aspirin Use		Expected	98.8%
Statin Use		Expected	88.8%
Tobacco-Free		Below Expected	81.4%
Optimal Vascular Care		Expected	55.5%
BP Control		Above Expected	85.0%
Daily Aspirin Use		Expected	84.9%
Statin Use		Expected	92.9%
Tobacco-Free		Expected	80.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).