

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	64.1%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Average	95.8%
Colorectal Cancer Screening		Expected	70.7%
Optimal Asthma Control - Adults		Expected	64.9%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	55.5%
HbA1c Control		Expected	81.1%
BP Control		Expected	88.4%
Daily Aspirin Use		Expected	98.8%
Statin Use		Expected	89.0%
Tobacco-Free		Expected	79.9%
Optimal Vascular Care		Expected	62.8%
BP Control		Above Expected	92.9%
Daily Aspirin Use		Expected	87.6%
Statin Use		Expected	97.4%
Tobacco-Free		Expected	81.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).