

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	65.3%
	Follow-up at Six Months	Above Expected	74.1%
	Response at Six Months	Above Expected	38.9%
	Remission at Six Months	Expected	20.4%
	Follow-up at 12 Months	Above Expected	74.1%
	Response at 12 Months	Expected	37.0%
	Remission at 12 Months	Above Expected	24.1%
Adolescent Mental Health and/or Depression Screening*		Average	98.3%
Colorectal Cancer Screening		Above Expected	76.2%
Optimal Asthma Control - Adults		Expected	64.2%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	44.0%
HbA1c Control		Expected	77.7%
BP Control		Expected	88.0%
Daily Aspirin Use		Expected	98.8%
Statin Use		Expected	91.6%
Tobacco-Free		Expected	77.7%
Optimal Vascular Care		Expected	55.3%
BP Control		Expected	81.2%
Daily Aspirin Use		Expected	89.4%
Statin Use		Expected	90.6%
Tobacco-Free		Expected	72.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).