

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	76.4%
	Follow-up at Six Months	Above Expected	77.1%
	Response at Six Months	Above Expected	44.3%
	Remission at Six Months	Above Expected	27.9%
	Follow-up at 12 Months	Above Expected	90.2%
	Response at 12 Months	Above Expected	47.5%
	Remission at 12 Months	Above Expected	34.4%
Adolescent Mental Health and/or Depression Screening*		Average	95.2%
Colorectal Cancer Screening		Below Expected	67.8%
Optimal Asthma Control - Adults		Expected	52.5%
Optimal Asthma Control - Children		Expected	52.9%
Optimal Diabetes Care		Expected	45.4%
HbA1c Control		Expected	72.0%
BP Control		Expected	85.6%
Daily Aspirin Use		Expected	99.5%
Statin Use		Expected	88.9%
Tobacco-Free		Below Expected	75.4%
Optimal Vascular Care		Expected	53.4%
BP Control		Expected	84.1%
Daily Aspirin Use		Expected	89.8%
Statin Use		Expected	92.1%
Tobacco-Free		Below Expected	73.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).