

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	40.5%
	Follow-up at Six Months	Above Expected	82.1%
	Response at Six Months	Expected	35.9%
	Remission at Six Months	Above Expected	25.6%
	Follow-up at 12 Months	Above Expected	74.4%
	Response at 12 Months	Above Expected	43.6%
	Remission at 12 Months	Expected	23.1%
Adolescent Mental Health and/or Depression Screening*		Above	100.0%
Colorectal Cancer Screening		Expected	65.5%
Optimal Asthma Control - Adults		Expected	51.7%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	51.3%
HbA1c Control		Expected	78.3%
BP Control		Expected	86.2%
Daily Aspirin Use		Expected	99.5%
Statin Use		Expected	91.0%
Tobacco-Free		Expected	78.8%
Optimal Vascular Care		Expected	50.6%
BP Control		Expected	87.9%
Daily Aspirin Use		Expected	90.1%
Statin Use		Expected	85.7%
Tobacco-Free		Expected	70.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).