

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	72.9%
	Follow-up at Six Months	Above Expected	77.1%
	Response at Six Months	Above Expected	39.6%
	Remission at Six Months	Above Expected	18.8%
	Follow-up at 12 Months	Above Expected	70.1%
	Response at 12 Months	Above Expected	38.2%
	Remission at 12 Months	Above Expected	21.5%
Adolescent Mental Health and/or Depression Screening*		Average	93.9%
Colorectal Cancer Screening		Expected	68.8%
Optimal Asthma Control - Adults		Above Expected	58.9%
Optimal Asthma Control - Children		Expected	57.1%
Optimal Diabetes Care		Above Expected	53.8%
HbA1c Control		Expected	74.9%
BP Control		Above Expected	96.1%
Daily Aspirin Use		Expected	99.3%
Statin Use		Above Expected	95.0%
Tobacco-Free		Below Expected	72.6%
Optimal Vascular Care		Above Expected	63.0%
BP Control		Above Expected	94.8%
Daily Aspirin Use		Above Expected	95.2%
Statin Use		Expected	91.0%
Tobacco-Free		Below Expected	70.2%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).