

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	74.6%
	Follow-up at Six Months	Expected	61.1%
	Response at Six Months	Above Expected	35.2%
	Remission at Six Months	Above Expected	29.6%
	Follow-up at 12 Months	Expected	61.1%
	Response at 12 Months	Above Expected	37.0%
	Remission at 12 Months	Above Expected	25.9%
Adolescent Mental Health and/or Depression Screening*		Average	100.0%
Colorectal Cancer Screening		Above Expected	75.6%
Optimal Asthma Control - Adults		Expected	65.2%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	46.5%
HbA1c Control		Expected	72.9%
BP Control		Above Expected	94.7%
Daily Aspirin Use		Expected	99.4%
Statin Use		Expected	92.9%
Tobacco-Free		Below Expected	70.6%
Optimal Vascular Care		Expected	58.0%
BP Control		Above Expected	93.8%
Daily Aspirin Use		Expected	90.2%
Statin Use		Below Expected	83.0%
Tobacco-Free		Expected	75.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).