

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	83.6%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		Above	100.0%
Colorectal Cancer Screening		Above Expected	83.8%
Optimal Asthma Control - Adults		Expected	71.9%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	59.5%
HbA1c Control		Expected	82.9%
BP Control		Expected	87.4%
Daily Aspirin Use		Expected	99.1%
Statin Use		Above Expected	95.5%
Tobacco-Free		Expected	83.8%
Optimal Vascular Care		Expected	55.9%
BP Control		Expected	86.4%
Daily Aspirin Use		Expected	91.5%
Statin Use		Expected	93.2%
Tobacco-Free		Expected	76.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).