

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	81.3%
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	74.9%
	Follow-up at Six Months	Above Expected	78.3%
	Response at Six Months	Above Expected	40.6%
	Remission at Six Months	Above Expected	22.6%
	Follow-up at 12 Months	Above Expected	67.9%
	Response at 12 Months	Above Expected	35.9%
	Remission at 12 Months	Above Expected	22.6%
Adolescent Mental Health and/or Depression Screening*		Above	97.5%
Colorectal Cancer Screening		Expected	73.7%
Optimal Asthma Control - Adults		Above Expected	70.0%
Optimal Asthma Control - Children		Above Expected	78.0%
Optimal Diabetes Care		Expected	51.5%
HbA1c Control		Expected	77.3%
BP Control		Above Expected	90.3%
Daily Aspirin Use		Expected	99.1%
Statin Use		Expected	87.7%
Tobacco-Free		Below Expected	77.7%
Optimal Vascular Care		Expected	55.3%
BP Control		Expected	85.5%
Daily Aspirin Use		Above Expected	95.3%
Statin Use		Expected	94.5%
Tobacco-Free		Below Expected	71.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).