

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	Average	91.7%
	Follow-up at Six Months	Expected	41.2%
	Response at Six Months	Expected	17.7%
	Remission at Six Months	Expected	8.8%
	Follow-up at 12 Months	Expected	47.1%
	Response at 12 Months	Expected	14.7%
	Remission at 12 Months	Expected	5.9%
Adult Depression	PHQ-9 Utilization*	Average	81.9%
	Follow-up at Six Months	Above Expected	74.1%
	Response at Six Months	Above Expected	31.3%
	Remission at Six Months	Above Expected	20.4%
	Follow-up at 12 Months	Above Expected	67.8%
	Response at 12 Months	Above Expected	35.9%
	Remission at 12 Months	Above Expected	23.0%
Adolescent Mental Health and/or Depression Screening*		Above	98.6%
Colorectal Cancer Screening		Above Expected	77.5%
Optimal Asthma Control - Adults		Above Expected	64.3%
Optimal Asthma Control - Children		Above Expected	67.4%
Optimal Diabetes Care		Above Expected	51.8%
HbA1c Control		Above Expected	80.7%
BP Control		Expected	80.0%
Daily Aspirin Use		Expected	99.3%
Statin Use		Above Expected	92.2%
Tobacco-Free		Expected	82.4%
Optimal Vascular Care		Expected	55.7%
BP Control		Expected	83.6%
Daily Aspirin Use		Expected	89.3%
Statin Use		Expected	94.0%
Tobacco-Free		Expected	78.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).