

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	79.8%
	Follow-up at Six Months	Above Expected	75.1%
	Response at Six Months	Above Expected	48.2%
	Remission at Six Months	Above Expected	31.8%
	Follow-up at 12 Months	Above Expected	67.2%
	Response at 12 Months	Above Expected	37.0%
	Remission at 12 Months	Above Expected	28.6%
Adolescent Mental Health and/or Depression Screening*		Above	98.8%
Colorectal Cancer Screening		Above Expected	76.2%
Optimal Asthma Control - Adults		Above Expected	69.7%
Optimal Asthma Control - Children		Above Expected	73.3%
Optimal Diabetes Care		Above Expected	58.1%
HbA1c Control		Expected	78.6%
BP Control		Above Expected	90.1%
Daily Aspirin Use		Expected	99.4%
Statin Use		Expected	92.1%
Tobacco-Free		Expected	85.3%
Optimal Vascular Care		Expected	61.0%
BP Control		Above Expected	88.1%
Daily Aspirin Use		Expected	90.6%
Statin Use		Expected	93.4%
Tobacco-Free		Expected	78.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).