

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	61.2%
	Follow-up at Six Months	Above Expected	80.8%
	Response at Six Months	Above Expected	38.5%
	Remission at Six Months	Above Expected	32.1%
	Follow-up at 12 Months	Above Expected	75.6%
	Response at 12 Months	Above Expected	39.7%
	Remission at 12 Months	Above Expected	25.6%
Adolescent Mental Health and/or Depression Screening*		Average	95.7%
Colorectal Cancer Screening		Above Expected	77.6%
Optimal Asthma Control - Adults		Expected	59.8%
Optimal Asthma Control - Children		Expected	61.9%
Optimal Diabetes Care		Expected	42.6%
HbA1c Control		Expected	68.3%
BP Control		Above Expected	87.8%
Daily Aspirin Use		Expected	99.0%
Statin Use		Expected	91.7%
Tobacco-Free		Below Expected	74.6%
Optimal Vascular Care		Expected	60.2%
BP Control		Expected	86.1%
Daily Aspirin Use		Above Expected	94.5%
Statin Use		Expected	94.2%
Tobacco-Free		Expected	74.8%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).