

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	46.1%
	Follow-up at Six Months	Expected	64.9%
	Response at Six Months	Expected	24.3%
	Remission at Six Months	Expected	12.2%
	Follow-up at 12 Months	Expected	59.5%
	Response at 12 Months	Expected	21.6%
	Remission at 12 Months	Expected	17.6%
Adolescent Mental Health and/or Depression Screening*		Below	70.4%
Colorectal Cancer Screening		Expected	74.8%
Optimal Asthma Control - Adults		Expected	61.0%
Optimal Asthma Control - Children		Expected	58.3%
Optimal Diabetes Care		Expected	54.4%
HbA1c Control		Expected	80.5%
BP Control		Above Expected	86.4%
Daily Aspirin Use		Expected	98.8%
Statin Use		Expected	92.5%
Tobacco-Free		Expected	82.3%
Optimal Vascular Care		Expected	57.3%
BP Control		Above Expected	87.8%
Daily Aspirin Use		Expected	88.2%
Statin Use		Expected	92.0%
Tobacco-Free		Below Expected	76.9%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).