

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	72.5%
	Follow-up at Six Months	Above Expected	69.8%
	Response at Six Months	Above Expected	36.5%
	Remission at Six Months	Above Expected	27.1%
	Follow-up at 12 Months	Expected	63.5%
	Response at 12 Months	Expected	25.0%
	Remission at 12 Months	Expected	13.5%
Adolescent Mental Health and/or Depression Screening*		Average	95.2%
Colorectal Cancer Screening		Above Expected	78.5%
Optimal Asthma Control - Adults		Expected	52.9%
Optimal Asthma Control - Children		Expected	47.0%
Optimal Diabetes Care		Expected	50.8%
HbA1c Control		Expected	78.5%
BP Control		Above Expected	85.4%
Daily Aspirin Use		Expected	99.1%
Statin Use		Expected	91.3%
Tobacco-Free		Below Expected	79.3%
Optimal Vascular Care		Expected	55.7%
BP Control		Expected	85.4%
Daily Aspirin Use		Above Expected	93.3%
Statin Use		Expected	93.3%
Tobacco-Free		Below Expected	73.1%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).