

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	65.7%
	Follow-up at Six Months	Above Expected	68.4%
	Response at Six Months	Expected	28.1%
	Remission at Six Months	Expected	14.0%
	Follow-up at 12 Months	Expected	57.9%
	Response at 12 Months	Expected	26.3%
	Remission at 12 Months	Expected	17.5%
Adolescent Mental Health and/or Depression Screening*		Above	99.3%
Colorectal Cancer Screening		Expected	72.1%
Optimal Asthma Control - Adults		Above Expected	61.1%
Optimal Asthma Control - Children		Expected	52.1%
Optimal Diabetes Care		Expected	51.6%
HbA1c Control		Expected	74.0%
BP Control		Above Expected	92.9%
Daily Aspirin Use		Expected	99.7%
Statin Use		Expected	90.9%
Tobacco-Free		Expected	79.6%
Optimal Vascular Care		Expected	60.7%
BP Control		Above Expected	90.6%
Daily Aspirin Use		Expected	92.0%
Statin Use		Expected	92.0%
Tobacco-Free		Expected	79.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).