

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	63.4%
	Follow-up at Six Months	Above Expected	76.9%
	Response at Six Months	Above Expected	29.1%
	Remission at Six Months	Above Expected	19.7%
	Follow-up at 12 Months	Above Expected	70.9%
	Response at 12 Months	Above Expected	28.2%
	Remission at 12 Months	Expected	15.4%
Adolescent Mental Health and/or Depression Screening*		Average	95.1%
Colorectal Cancer Screening		Expected	67.5%
Optimal Asthma Control - Adults		Above Expected	67.2%
Optimal Asthma Control - Children		Expected	61.7%
Optimal Diabetes Care		Expected	49.8%
HbA1c Control		Expected	77.1%
BP Control		Above Expected	87.7%
Daily Aspirin Use		Expected	99.4%
Statin Use		Expected	88.3%
Tobacco-Free		Below Expected	76.8%
Optimal Vascular Care		Expected	60.0%
BP Control		Above Expected	87.8%
Daily Aspirin Use		Above Expected	93.6%
Statin Use		Expected	91.9%
Tobacco-Free		Expected	75.3%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).