

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Average	77.9%
	Follow-up at Six Months	Above Expected	76.5%
	Response at Six Months	Above Expected	31.1%
	Remission at Six Months	Expected	17.7%
	Follow-up at 12 Months	Above Expected	74.0%
	Response at 12 Months	Above Expected	32.8%
	Remission at 12 Months	Above Expected	24.4%
Adolescent Mental Health and/or Depression Screening*		Average	91.9%
Colorectal Cancer Screening		Above Expected	75.3%
Optimal Asthma Control - Adults		Above Expected	64.0%
Optimal Asthma Control - Children		Expected	70.0%
Optimal Diabetes Care		Above Expected	56.1%
HbA1c Control		Above Expected	81.1%
BP Control		Above Expected	89.7%
Daily Aspirin Use		Expected	99.3%
Statin Use		Expected	90.8%
Tobacco-Free		Expected	82.0%
Optimal Vascular Care		Above Expected	67.1%
BP Control		Above Expected	87.7%
Daily Aspirin Use		Above Expected	93.3%
Statin Use		Expected	95.2%
Tobacco-Free		Expected	81.4%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).