

## Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Below	20.4%
	Follow-up at Six Months	Expected	49.5%
	Response at Six Months	Expected	11.2%
	Remission at Six Months	Expected	5.6%
	Follow-up at 12 Months	Expected	39.3%
	Response at 12 Months	Expected	10.3%
	Remission at 12 Months	Below Expected	2.8%
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		-	-
Optimal Asthma Control - Adults		-	-
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		-	-
HbA1c Control		-	-
BP Control		-	-
Daily Aspirin Use		-	-
Statin Use		-	-
Tobacco-Free		-	-
Optimal Vascular Care		-	-
BP Control		-	-
Daily Aspirin Use		-	-
Statin Use		-	-
Tobacco-Free		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

\*Measure not risk adjusted

### Ratings

**Above:** Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

**Average:** Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

**Below:** Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).