

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adolescent Mental Health and/or Depression Screening*		-	-
Colorectal Cancer Screening		-	-
Optimal Asthma Control - Adults		-	-
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Expected	44.2%
<i>HbA1c Control</i>		<i>Expected</i>	<i>68.6%</i>
<i>BP Control</i>		<i>Expected</i>	<i>83.9%</i>
<i>Daily Aspirin Use</i>		<i>Expected</i>	<i>98.3%</i>
<i>Statin Use</i>		<i>Expected</i>	<i>83.3%</i>
<i>Tobacco-Free</i>		<i>Above Expected</i>	<i>91.5%</i>
Optimal Vascular Care		-	-
<i>BP Control</i>		-	-
<i>Daily Aspirin Use</i>		-	-
<i>Statin Use</i>		-	-
<i>Tobacco-Free</i>		-	-

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).