

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Expected	45.4%
	Response at Six Months	Expected	10.3%
	Remission at Six Months	Expected	5.2%
	Follow-up at 12 Months	Expected	39.2%
	Response at 12 Months	Below Expected	6.2%
	Remission at 12 Months	Below Expected	4.1%
Adolescent Mental Health and/or Depression Screening*		Average	95.0%
Colorectal Cancer Screening		Above Expected	81.7%
Optimal Asthma Control - Adults		Expected	56.7%
Optimal Asthma Control - Children		Expected	74.3%
Optimal Diabetes Care		Expected	55.6%
HbA1c Control		Expected	81.7%
BP Control		Expected	83.7%
Daily Aspirin Use		Expected	99.3%
Statin Use		Above Expected	93.2%
Tobacco-Free		Expected	84.5%
Optimal Vascular Care		Expected	59.2%
BP Control		Expected	83.7%
Daily Aspirin Use		Expected	91.9%
Statin Use		Expected	94.0%
Tobacco-Free		Expected	79.0%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).