

Clinical Quality Measures

2023 measurement year

Measure		Rating	Rate
Adolescent Depression	PHQ-9 Utilization*	-	-
	Follow-up at Six Months	-	-
	Response at Six Months	-	-
	Remission at Six Months	-	-
	Follow-up at 12 Months	-	-
	Response at 12 Months	-	-
	Remission at 12 Months	-	-
Adult Depression	PHQ-9 Utilization*	Above	100.0%
	Follow-up at Six Months	Below Expected	38.5%
	Response at Six Months	Expected	15.6%
	Remission at Six Months	Expected	10.4%
	Follow-up at 12 Months	Expected	41.7%
	Response at 12 Months	Expected	20.8%
	Remission at 12 Months	Expected	11.5%
Adolescent Mental Health and/or Depression Screening*		Above	98.9%
Colorectal Cancer Screening		Above Expected	80.1%
Optimal Asthma Control - Adults		Above Expected	69.2%
Optimal Asthma Control - Children		-	-
Optimal Diabetes Care		Above Expected	58.9%
HbA1c Control		Expected	82.5%
BP Control		Expected	84.1%
Daily Aspirin Use		Expected	96.5%
Statin Use		Above Expected	95.9%
Tobacco-Free		Expected	85.0%
Optimal Vascular Care		Expected	53.1%
BP Control		Expected	79.6%
Daily Aspirin Use		Expected	82.6%
Statin Use		Expected	92.0%
Tobacco-Free		Expected	81.5%

- Clinic not assigned to measure, did not have results, or did not meet reporting threshold

*Measure not risk adjusted

Ratings

Above: Clinic's actual rate is significantly above its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted measures).

Average: Clinic's actual rate is not statistically different than its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).

Below: Clinic's actual rate is significantly below its expected rate (for risk adjusted measures) or the statewide rate (for unadjusted).